Untitled Document Page 1 of 3

Musca-Cide™ FLY BAIT

ATTRACTS AND KILLS MORE FEMALE FLIES

- Advanced Bait Formula Contains TRICOLURETM Sex Attractant and Feeding Adjuvants
- Attracts and Kills House Flies, Plus Many Other Fly Species Listed on This Label
- Attracts Both Sexes of Flies to the Bait
- ∠ Kills Quickly

ACTIVE INGREDIENTS:

Methomyl [S-Methyl N-[(methylcarbamoyl) oxy] thioacetimidate] 1.000% Z-9 Tricosene 0.025% INERT INGREDIENTS: 98.975%

Keep Out Of Reach Of Children CAUTION See back panel for additional cautions NET WT. 11.34 kg (25 LBS.)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Do not contaminate water, feed or foodstuffs by use, storage or disposal. Do not apply where poultry, animals, pets or children can contact bait. Do not use in homes or where milk is processed or stored. Do not feed treated garbage. Do not use in serving areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

STATEMENT OF PRACTICAL TREATMENT

IF INGESTED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician immediately.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Wash contaminated clothing before reuse.

IF IN EYES: Flush with plenty of water, call a physician immediately.

NOTE TO PHYSICIAN: Methomyl is a cholinesterase inhibitor. Atropine is antidotal. Administer intravenously or intramuscularly if IV injection is not possible.

ENVIRONMENTAL HAZARDS

Exposed surfaces treated with this bait are hazardous to birds and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters.

Musca-Cide is an advanced formula bait containing Tricolure sex attractant and feeding adjuvants which attract both sexes of the house fly and many other pest fly species including

Untitled Document Page 2 of 3

little house flies, flesh flies, blow flies, bazaar flies (Musca sorbens) and eye gnats directly to the bait.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

NOTE: This product may not be used (as either a scatter bait or in a bait station) around or inside homes or any other places where children are likely to be present. Residential use of this product is prohibited.

For use as a Scatter Bait: Use of Musca-Cide Bait as a scatter bait is limited to certain commercial agricultural production areas where children and pets are not likely to be present. Use only around the outside of feed lots, dairies, broiler houses, hog houses, livestock barns, meat and poultry processing establishments, beverage plants, canneries and food processing establishments. Bait may be used on walkways in caged layer houses.

Bait should be applied to specified areas of high fly population daily or as needed to reduce fly populations. Scatter bait from container or other device at the rate of approximately 4 ounces per 500 square feed of fly feeding surface. Do not allow food producing animals to have access to treated areas.

For Use in the Bait Station: For stables and other commercial (non-residential) use sites not specified above, Musca-Cide Bait must be used in a bait

station. Fill bait chamber to bottom of hopper by pouring 2 ounces of bait into hopper. Use at the rate of approximately 2 bait stations per 500 square feet of fly feeding area. Bait stations must be inaccessible to children and food producing animals. Hang bait stations out of reach of children and food producing animals at a height of at least 4 feet above the ground.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

EPA EST. NO.: 39578-TX-1
EPA REG. NO.: 270-255
U.S. PATENT NO. 5,008,107
Manufactured For:
FARNAM COMPANIES, INC.
Pest Control Division
P.O. Box 34820, Phoenix, Arizona 85067-4820

©1999 Farnam Companies, Inc

Product Code: 056226

Untitled Document Page 3 of 3

Printed in U.S.A. 9EE9