

PULL HERE TO OPEN ►

talon®-G

Rodenticide Mini-Pellets



For Control of Commensal Rats and House Mice

Second Generation Anticoagulant

Kills Warfarin-Resistant Norway Rats and House Mice

Rodents may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

Active Ingredient:

Brodifacoum 0.005%

Other Ingredients: 99.995%

Total: 100.000%

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use on booklet.

EPA Reg. No. 100-1057

EPA Est. 61282-WI-1

Product of the United Kingdom

Formulated in the USA

SCP 1057A-L1C 1105

For use of optional measuring cone, see instructions under Application Directions.

5 pounds (3/32" mini-pellets)

Net Weight

syngenta

Talon®-G

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May be harmful or fatal if swallowed. Keep away from humans, domestic animals and pets. Wash hands after handling bait.

IF BAIT IS EATEN BY ANIMALS OR PETS, CALL A VETERINARIAN AT ONCE.

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
If swallowed	<ul style="list-style-type: none">• Call a poison control center, doctor, or HOT LINE NUMBER immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	
NOTE TO PHYSICIAN OR VETERINARIAN This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K1 are indicated, as in poisoning from overdose of bishydroxy coumarin. For human cases, vitamin K1 is antidotal at doses of 10-20 mg (not mg/kg). For animal cases, vitamin K1 is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to 2 weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.	

Environmental Hazards

This product is toxic to fish, birds, and wildlife. This product can pose a secondary hazards to birds of prey and mammals. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children or pets.

Talon®-G

2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under 6 years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

Use Restrictions

For control of Norway rats, Roof rats, and House mice in and around homes, industrial, commercial, agricultural and public buildings, and similar manmade structures. This product may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

Selection of Treatment Areas

Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS

Norway and Roof Rats

Apply 4–16 oz. of bait (usually at intervals of 15–30 ft.) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

House Mice

Apply 1/4 to 1/2 ounce of bait per placement. Space placements at intervals of 8–12 ft. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

Follow-up

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead rodents and left-over bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. Where a continuous source of infestation is present, maintain permanent bait stations and replenish with census or toxic bait as needed.

Use of Measuring Cone

For use of the Talon-G Pellet measuring cone, please refer to the supplemental sheet **Talon-G Pellet Measuring Cone Instructions**, as packed with this container, or available from your Distributor, or by calling Syngenta Crop Protection, Inc. at 1-800-344-9481.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage

Store only in original container in dry place inaccessible to children and pets.

Pesticide Disposal

Dispose of wastes resulting from the use of this product at an approved waste disposal facility or call your local solid waste agency for alternative disposal instructions. Never place unused product down any indoor or outdoor drain.

Pesticide Container

Do not reuse empty container except for holding additional Talon®-G Rodenticide Mini-Pellets Contains Bitrex®. Dispose of in a sanitary landfill or by other procedures allowed by State and local authorities.

Talon® and the Syngenta logo are trademarks of a Syngenta Group Company.

Bitrex® is a registered trademark of Macfarlan Smith Ltd., Edinburgh, Scotland.

©2006 Syngenta

For non-emergency (e.g. current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 1057A-L1C 1105

talon®-G

KEEP OUT OF REACH OF CHILDREN. CAUTION

See additional precautionary statements and directions for use on booklet.

Rodenticide Mini-Pellets



For Control of Commensal
Rats and House Mice

Second Generation Anticoagulant

Kills Warfarin-Resistant Norway Rats and
House Mice

Rodents may consume a lethal dose in one
feeding with first dead rodents appearing
four or five days after treatment begins.

Active Ingredient:	
Brodifacoum	0.005%
Other Ingredients:	99.995%
Total:	100.000%

EPA Reg. No. 100-1057
EPA Est. 61282-WI-1

Talon® and the Syngenta logo are trade-
marks of a Syngenta Group Company.
Bitrex® is a registered trademark of
Macfarlan Smith Ltd., Edinburgh, Scotland.

©2006 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 1057A-L1C 1105

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May be harmful or fatal if swallowed. Keep away from humans, domestic
animals and pets. Wash hands after handling bait.

IF BAIT IS EATEN BY ANIMALS OR PETS, CALL A VETERINARIAN AT ONCE.

FIRST AID

Have the product container or label with you when calling a poison control
center or doctor, or going for treatment.

If swallowed: Call a poison control center, doctor, or **HOT LINE NUMBER**
immediately for treatment advice. Have person sip a glass of water if able to
swallow. Do not induce vomiting unless told to do so by the poison control
center or doctor.

HOT LINE NUMBER: For 24-Hour Medical Emergency Assistance (Human or
Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call
1-800-888-8372.

NOTE TO PHYSICIAN OR VETERINARIAN: This product may reduce the
clotting ability of the blood and cause hemorrhaging. If poisoning occurs,
intramuscular and oral administration of vitamin K₁ are indicated, as in
poisoning from overdose of bishydroxy coumarin. For human cases, vitamin K₁
is antidotal at doses of 10-20 mg (**not** mg/kg). For animal cases, vitamin K₁ is
antidotal at 2-5 mg/kg. Repeated doses may need to be given up to 2 weeks
(based on monitoring of prothrombin times). In severe cases, blood transfu-
sions may be necessary.

Environmental Hazards

This product is toxic to fish, birds, and wildlife. This product can pose a
secondary hazards to birds of prey and mammals. Do not apply directly to
water, or to areas where surface water is present or to intertidal areas below
the mean high water mark.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage

Store only in original container in dry place inaccessible to children and pets.

Pesticide Disposal

Dispose of wastes resulting from the use of this product at an approved waste
disposal facility or call your local solid waste agency for alternative disposal
instructions. Never place unused product down any indoor or outdoor drain.

Pesticide Container

Do not reuse empty container except for holding additional Talon®-G
Rodenticide Mini-Pellets Contains Bitrex®. Dispose of in a sanitary landfill or
by other procedures allowed by State and local authorities.

5 pounds (3/32" mini-pellets)

Net Weight

